

Problem sheet 2

Due date: Oct. 23, 2018.

Problem 5

Let $\varphi: A \rightarrow B$ be a ring homomorphism, and let $f: \text{Spec } B \rightarrow \text{Spec } A$ denote the map attached to φ .

(a) Let $\mathfrak{b} \subseteq B$ be an ideal. Prove that

$$\overline{f(V(\mathfrak{b}))} = V(\varphi^{-1}(\mathfrak{b}))$$

(where $\bar{}$ denotes closure).

(b) Assume that φ is surjective. Prove that f induces a homeomorphism from $\text{Spec } B$ onto $V(\ker(\varphi))$.

(c) Prove that the image of f is dense in $\text{Spec } A$ if and only if every element of $\ker(\varphi)$ is nilpotent.

Problem 6

Let A be a ring, $f \in A$ and $X := D(f) \subseteq \text{Spec } A$ with the subspace topology. Prove that the topological space X is quasi-compact (i.e., for every cover $X = \bigcup_{i \in I} U_i$ by open subsets U_i , there exists a finite subset $I' \subseteq I$ with $X = \bigcup_{i \in I'} U_i$).

Problem 7

Let (I, \leq) be a partially ordered set. An *inductive system* of sets (with index set I) is a family X_i , $i \in I$, of sets together with maps $\varphi_{ji}: X_i \rightarrow X_j$ for all pairs $i, j \in I$ with $i \leq j$, such that

$$\varphi_{ii} = \text{id}_{X_i}, \quad \varphi_{kj} \circ \varphi_{ji} = \varphi_{ki}$$

for all $i \leq j \leq k \in I$.

A set C together with maps $\psi_i: X_i \rightarrow C$ such that $\psi_j \circ \varphi_{ji} = \psi_i$ for all $i \leq j$ is called a *colimit* (or *direct limit* or *inductive limit*) if it satisfies the following “universal property”: For every set T together with maps $\xi_i: X_i \rightarrow T$ such that $\xi_j \circ \varphi_{ji} = \xi_i$ for all $i \leq j$, there exists a unique map $\chi: C \rightarrow T$ such that $\chi \circ \psi_i = \xi_i$ for all i . The colimit is also denoted by $\text{colim}_{i \in I} X_i$ or by $\varinjlim_{i \in I} X_i$. (The maps φ_{ij} and ψ_i are usually omitted from the notation.)

- (a) Suppose that I is *directed*, i.e., for all $i, j \in I$ there exists $k \in I$ with $i \leq k$ and $j \leq k$. Let (X_i, φ_{ji}) be an inductive system of sets, let U be the disjoint union

$$U = \coprod_{i \in I} X_i,$$

and consider the following relation on U : for $x, y \in U$, say $x \in X_i, y \in X_j$, we set $x \sim y$ if and only if there exists $k \geq i, j$ with $\varphi_{ki}(x) = \varphi_{kj}(y)$. Prove that \sim is an equivalence relation and that the set U/\sim of equivalence classes together with the natural maps $X_i \rightarrow U/\sim$ is a colimit of the system (X_i, φ_{ji}) .

- (b) Assume that I is directed, that all X_i are subsets of a set X , that $i \leq j$ if and only if $X_i \subseteq X_j$, and that the maps φ_{ji} are the inclusion maps. Prove that the union $C := \bigcup_i X_i$ with the inclusion maps $X_i \rightarrow C$ is a colimit of the X_i .

Problem 8

Let I be a partially ordered set.

- (a) Define the notion of *colimit* (with index set I) in an arbitrary category.
- (b) Suppose that I is directed. Prove that all colimits with index set I in the category of abelian groups exist.
- (c) Suppose that I is directed. Prove that the functor colim_i on the category of abelian groups is exact, i.e.: Let $(A_i, \varphi_{ji}), (B_i, \psi_{ji}), (C_i, \xi_{ji})$ be inductive systems of abelian groups indexed by I . Suppose that for each i , there are short exact sequences

$$0 \rightarrow A_i \rightarrow B_i \rightarrow C_i \rightarrow 0$$

which are compatible with the maps $\varphi_{ji}, \psi_{ji}, \xi_{ji}$ in the obvious sense. Prove that these sequences induce a sequence

$$0 \rightarrow \text{colim}_i A_i \rightarrow \text{colim}_i B_i \rightarrow \text{colim}_i C_i \rightarrow 0$$

which is again exact.